

Renormalization of the Cabibbo-Kobayashi-Maskawa Matrix at One-Loop Level

Yong Zhou^a

^a *Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China*

We have investigated the present renormalization prescriptions of Cabibbo-Kobayashi-Maskawa (CKM) matrix at one-loop level. We emphasize at one prescription which is formulated with reference to the case of no mixing of quark's generations and point out that it doesn't make the physical amplitude involving quark mixing ultraviolet finite. We then propose a revised prescription to solve this problem and simultaneously keep the unitarity of the CKM matrix. Through explicit calculations we also prove that in order to keep the CKM matrix gauge independent the unitarity of the CKM matrix must be preserved.

11.10.Gh, 12.15.Lk, 12.15.Hh

I. INTRODUCTION

Since the exact examination of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [1–3] has been developed quickly, the renormalization of CKM matrix becomes very important [4]. This was realized for the Cabibbo angle with two fermion generations by Marciano and Sirlin [5] and for the CKM matrix of the three-generation SM by Denner and Sack [6] more than a decade ago. In recent years many people have discussed this issue [7], but a completely self-consistent scheme has been not obtained. In this paper we will study one of the present prescriptions and give some instructive conclusions.

In general, a CKM matrix renormalization prescription should satisfy the three criterions [8]:

1. In order to let the transition amplitude of any physical process involving quark mixing ultraviolet finite, the CKM counterterm must cancel out the ultraviolet divergence left in the loop-corrected amplitudes.
2. It must guarantee such transition amplitude gauge parameter independent [9], which is a fundamental requirement.
3. SM requires the bare CKM matrix V^0 is unitary,

$$\sum_k V_{ik}^0 V_{jk}^{0*} = \delta_{ij} \quad (1)$$

with i, j, k the generation index and δ_{ij} the unit matrix element. If we split the bare CKM matrix element into the renormalized one and its counterterm

$$V_{ij}^0 = V_{ij} + \delta V_{ij} \quad (2)$$

and keep the unitarity of the renormalized CKM matrix, the unitarity of the bare CKM matrix at one-loop level requires

$$\sum_k (\delta V_{ik} V_{jk}^* + V_{ik} \delta V_{jk}^*) = 0 \quad (3)$$

Until now there are many papers discussing this problem. The modified minimal subtraction ($\overline{MS}$) scheme [10,11] is the simplest one, but it introduces the μ^2 -dependent terms which are very complicated to be dealt with. In the on-shell renormalization framework, the early prescription [6] used the $SU_L(2)$ symmetry of SM to relate the CKM counterterm with the quark's wave-function renormalization constants (WRC) [12]. Although it is a delicate and simple prescription, it reduces the physical amplitude involving quark mixing gauge dependent¹ [13–15]. A revised prescription is to replace the on-shell quark WRC in the CKM counterterms with the ones calculated at zero

¹This is easy to be understood since the $SU_L(2)$ symmetry of SM has been broken by the Higgs mechanism

momentum [13]. Another revised prescription [15] is to rearrange the off-diagonal quark WRC in a manner similar to the pinch technique [16].

Different from the idea of Ref. [6], another idea is to formulate the CKM renormalization prescription with reference to the case of no mixing of quark's generations. This has been done in Ref. [17,8] at one-loop level. In the following section we will introduce this prescription and discuss the defect of the prescription in Ref. [17]. The revised prescription will be given in the following section. In section 4 we will discuss the relationship between the unitarity and the gauge independence of the CKM matrix through explicit calculations. Lastly we give our conclusions.

II. BARROSO'S CKM MATRIX RENORMALIZATION PRESCRIPTION

The main idea of Barroso's prescription is to renormalize the transition amplitude of W gauge boson decaying into two quarks equal to the same amplitude but in the case of no mixing of quark's generations. In order to elaborate this idea clearly we firstly introduce the one-loop decaying amplitude of $W^+ \rightarrow u_i \bar{d}_j$ [17]

$$T_1 = A_L [V_{ij}(F_L + \frac{\delta g}{g} + \frac{1}{2}\delta Z_W + \frac{1}{2}\delta \bar{Z}_{ii}^{uL} + \frac{1}{2}\delta Z_{jj}^{dL}) + \sum_{k \neq i} \frac{1}{2}\delta \bar{Z}_{ik}^{uL} V_{kj} + \sum_{k \neq j} \frac{1}{2} V_{ik} \delta Z_{kj}^{dL} + \delta V_{ij}] + V_{ij} [A_R F_R + B_L G_L + B_R G_R], \quad (4)$$

with g and δg the $SU(2)$ coupling constant and its counterterm, δZ_W the W boson WRC, $\delta \bar{Z}^{uL}$ and δZ^{dL} the left-handed up-type and down-type quark's WRC [7,18], and

$$A_L = \frac{g}{\sqrt{2}} \bar{u}_i(p_1) \not{\epsilon} \gamma_L \nu_j(q - p_1), \\ B_L = \frac{g}{\sqrt{2}} \bar{u}_i(p_1) \frac{\epsilon \cdot p_1}{M_W} \gamma_L \nu_j(q - p_1). \quad (5)$$

with ϵ^μ the W boson polarization vector, γ_L and γ_R the left-handed and right-handed chiral operators, and M_W the W boson mass. Similarly, replacing γ_L with γ_R in the equations we can define A_R and B_R respectively. $F_{L,R}$ and $G_{L,R}$ are four form factors. The main idea of Ref. [17] is to choose the CKM counterterm to let the amplitude T_1 similar as the amplitude of W boson decaying into two leptons. As we know if there is no mixing of lepton's generations the amplitude of $W^+ \rightarrow \nu_i \bar{e}_i$ is gauge independent and ultraviolet finite after introducing proper physical parameter's counterterms, except for CKM counterterm. So if the amplitude of $W^+ \rightarrow u_i \bar{d}_j$ is renormalized equal to the same amplitude but in the case of no mixing of quark's generations, we will get a gauge independent and ultraviolet finite amplitude. This procedure will determine the CKM counterterm. The key to this problem is to find the amplitude of $W^+ \rightarrow u_i \bar{d}_j$ in the case of no mixing of quark's generations. Barroso suggested that such amplitude should be the following form at one-loop level [17]:

$$T_1 = V_{ij} [A_L(F_L + \frac{\delta g}{g} + \frac{1}{2}\delta Z_W + \frac{1}{2}\delta \bar{Z}_{ii[l]}^{uL} + \frac{1}{2}\delta Z_{jj[l]}^{dL}) + A_R F_R + B_L G_L + B_R G_R] \quad (6)$$

where the subscript "[l]" denotes the quantity is obtained by replacing CKM matrix elements with unit matrix elements. It is easy to obtain [17]

$$\delta V_{ij} = -\frac{1}{2} \sum_k [\delta \bar{Z}_{ik}^{uL} V_{kj} + V_{ik} \delta Z_{kj}^{dL}] + \frac{1}{2} V_{ij} [\delta \bar{Z}_{ii[l]}^{uL} + \delta Z_{jj[l]}^{dL}] \quad (7)$$

But in fact such CKM counterterms don't make the decaying amplitude T_1 ultraviolet finite when $i \neq j$. It is easy to calculate the ultraviolet part of T_1 using δV as CKM counterterm

$$T_1|_{UV-divergence} = \frac{\alpha V_{ij} \Delta}{32\pi M_W^2 s_W^2} (m_{d,i}^2 - m_{d,j}^2 + m_{u,j}^2 - m_{u,i}^2)$$

with α the fine structure constant, s_W the sine of the weak mixing angle θ_W , $m_{d,i}$ and $m_{d,j}$ the down-type quark's masses, $m_{u,i}$ and $m_{u,j}$ the up-type quark's masses, and $\Delta = 2/(D-4) + \gamma_E - \ln(4\pi) + \ln(M_W^2/\mu^2)$ (D is the space-time dimensionality, γ_E is the Euler's constant and μ is an arbitrary mass parameter). This result shows that when $i \neq j$ the decaying amplitude of $W^+ \rightarrow u_i \bar{d}_j$ is ultraviolet divergent.

III. ONE-LOOP RENORMALIZATION OF CKM MATRIX

We argue that such mistake comes from the one-sided knowledge about the difference between the two cases of having and not having mixing of quark's generations. In the case of no mixing of quark's generations only the quarks in one generation can appear at a fermion line in a Feynman diagram, so only the quarks which are in the same generations as the external-line quarks can appear at the amplitude of $W^+ \rightarrow u_i \bar{d}_j$ in the case of no mixing of quark's generations, except for the counterterms δg and δZ_W . That's to say only two kinds of quarks: u_i and d_j are present in the renormalized amplitude of $W^+ \rightarrow u_i \bar{d}_j$, except for δg and δZ_W because they have nothing to do with fermion. This point has been implicated in Ref. [8]. Thus different from Eq.(6), the amplitude T_1 will be renormalized as follows

$$T_1 = V_{ij}[A_L(F_L + \frac{\delta g}{g} + \frac{1}{2}\delta Z_W + \frac{1}{2}\delta \bar{Z}_{ii}^{uL} m_{d,i} \rightarrow m_{d,j} + \frac{1}{2}\delta Z_{jj}^{dL} m_{u,j} \rightarrow m_{u,i}) + A_R F_R + B_L G_L + B_R G_R] \quad (8)$$

So the CKM counterterm is obtained compared with Eq.(4) and Eq.(8)

$$\delta V_{ij} = -\frac{1}{2} \sum_k [\delta \bar{Z}_{ik}^{uL} V_{kj} + V_{ik} \delta Z_{kj}^{dL}] + \frac{1}{2} V_{ij} [\delta \bar{Z}_{ii}^{uL} m_{d,i} \rightarrow m_{d,j} + \delta Z_{jj}^{dL} m_{u,j} \rightarrow m_{u,i}] \quad (9)$$

Our calculations have proven that this CKM counterterm is gauge independent and makes the physical amplitude T_1 ultraviolet convergent.

Ref. [8] has pointed out that such a CKM counterterm maybe not satisfy the unitary condition of Eq.(3), but an explicit Proof has not been given. Here we will do an analytical calculation to show in what degree δV satisfies the unitary condition. At one-loop level only four diagrams need to be considered when we calculate the CKM counterterm in Eq.(9), as shown in Fig.1.




FIG. 1. Quark's self-energy diagrams that contribute to the CKM counterterm in Eq.(10).

We have used the software packages *FeynArts* [19] to draw the Feynman diagrams and generate the corresponding Feynman amplitudes, then used *FeynCalc* [20] to calculate these Feynman amplitudes. It is easy to get the analytical result of δV_{ij} since the quark's self-energy functions at one-loop level are simple. In order to check the unitary condition of Eq.(3) analytically, we can use the Taylor's series, $(m_{quark}^2/M_W^2)^n$, to expand δV_{ij} . The one- and two-order results are shown as follows:

$$\delta V_{ij}^{(1)} = \frac{\alpha(6\Delta-11)}{128\pi M_W^2 s_W^2} \left[-\frac{2 \sum_{k,l \neq j} m_{d,j} m_{u,k}^2 V_{il} V_{kl}^* V_{kj}}{m_{d,l} - m_{d,j}} + \frac{2 \sum_{k,l} m_{d,j} m_{u,k}^2 V_{il} V_{kl}^* V_{kj}}{m_{d,l} + m_{d,j}} - \frac{2 \sum_{k \neq i,l} m_{u,i} m_{d,l}^2 V_{il} V_{kl}^* V_{kj}}{m_{u,k} - m_{u,i}} + \frac{2 \sum_{k,l} m_{u,i} m_{d,l}^2 V_{il} V_{kl}^* V_{kj}}{m_{u,k} + m_{u,i}} \right] + V_{ij} (\sum_k V_{ik} V_{ik}^* m_{d,k}^2 + \sum_k V_{kj} V_{kj}^* m_{u,k}^2 - 2m_{d,j}^2 - 2m_{u,i}^2). \quad (10)$$

$$\begin{aligned} \delta V_{ij}^{(2)} = & \frac{\alpha}{64\pi M_W^4 s_W^2} [V_{ij} (12 \ln \frac{m_{d,j}^2}{M_W^2} m_{d,j}^4 + 6m_{d,j}^4 + 12 \ln \frac{m_{u,i}^2}{M_W^2} m_{u,i}^4 + 6m_{u,i}^4 + 32m_{d,j}^2 m_{u,i}^2 - \\ & \sum_k V_{ik} V_{ik}^* (6 \ln \frac{m_{d,k}^2}{M_W^2} m_{d,k}^4 + 3m_{d,k}^4 + 12m_{u,i}^2 m_{d,k}^2) - \sum_k V_{kj} V_{kj}^* (6 \ln \frac{m_{u,k}^2}{M_W^2} m_{u,k}^4 + 3m_{u,k}^4 + 12m_{d,j}^2 m_{u,k}^2)) \\ & + \frac{2}{m_{d,l} - m_{d,j}} \sum_{k,l \neq j} m_{d,j} m_{u,k}^2 (4m_{d,j}^2 + 6 \ln \frac{m_{u,k}^2}{M_W^2} m_{u,k}^2 + 3m_{u,k}^2) V_{il} V_{kl}^* V_{kj} \\ & - \frac{2}{m_{d,l} + m_{d,j}} \sum_{k,l} m_{d,j} m_{u,k}^2 (4m_{d,j}^2 + 6 \ln \frac{m_{u,k}^2}{M_W^2} m_{u,k}^2 + 3m_{u,k}^2) V_{il} V_{kl}^* V_{kj} \\ & + \frac{2}{m_{u,k} - m_{u,i}} \sum_{k \neq i,l} m_{u,i} m_{d,l}^2 (4m_{u,i}^2 + 6 \ln \frac{m_{d,l}^2}{M_W^2} m_{d,l}^2 + 3m_{d,l}^2) V_{il} V_{kl}^* V_{kj} \\ & - \frac{2}{m_{u,k} + m_{u,i}} \sum_{k,l} m_{u,i} m_{d,l}^2 (4m_{u,i}^2 + 6 \ln \frac{m_{d,l}^2}{M_W^2} m_{d,l}^2 + 3m_{d,l}^2) V_{il} V_{kl}^* V_{kj}]. \end{aligned} \quad (11)$$

where the superscript "(1)" and "(2)" denote the one-order and two-order results of δV_{ij} about the series m_{quark}^2/M_W^2 . The R_ξ -gauge [21] has been used. Replacing δV with $\delta V^{(1)} + \delta V^{(2)}$ in Eq.(3), we find at one-loop level it satisfies the unitary condition.

But when we consider the three-order result of δV about the series m_{quark}^2/M_W^2 , we find it doesn't satisfy the unitary condition, as shown below:

$$\sum_k (\delta V_{ki}^{(3)*} V_{kj} + V_{ki}^* \delta V_{kj}^{(3)}) = \frac{9\alpha}{128\pi M_W^6 s_W^2} [\sum_k m_{u,k}^2 (m_{d,i}^4 - 2m_{d,j}^2 m_{d,i}^2 + m_{d,j}^4 + m_{d,i}^2 m_{u,k}^2 + m_{d,j}^2 m_{u,k}^2) V_{ki}^* V_{kj} - 2 \sum_{k,l,n} m_{u,k}^2 m_{d,l}^2 m_{u,n}^2 V_{ki}^* V_{kl} V_{nl}^* V_{nj}] \neq 0 \quad (12)$$

This result shows that δV doesn't comply with the unitary criterion. But from this result we can see that the deviation of δV from the unitary condition is very small since the quark's masses are very small compared with M_W , except for m_t . Calculating to five-order result of δV about the series m_{quark}^2/M_W^2 , we find the largest deviation of $\delta V^\dagger V + V^\dagger \delta V$ from 0 is proportional to $\alpha |V_{3i \neq 3}| m_b^2 m_t^8 / (s_W^2 M_W^{10}) \sim 10^{-7}$, which is very small compared with the present measurement precision of the CKM matrix elements. Thus in actual calculations we can use Eq.(9) as the definition of the CKM counterterm.

Of course an severe CKM renormalization prescription should satisfy the unitary condition severely. Diener and Kniehl have proposed a prescription which can shift a proper CKM counterterm which has right ultraviolet divergence and gauge independent to satisfy the three criterions proposed in Ref. [8], as shown below [8]

$$\delta \bar{V}_{ij} = \frac{1}{2} (\delta V_{ij} - \sum_{k,l} V_{ik} \delta V_{kl}^\dagger V_{lj}) \quad (13)$$

It is easy to prove that $\delta \bar{V}$ satisfies the unitary condition of Eq.(3). Ref. [8] has pointed out that if the ultraviolet divergence of δV satisfies the unitary condition of Eq.(3) $\delta \bar{V}$ will has the same ultraviolet divergence as δV . Because we have checked that the δV in Eq.(9) is gauge independent and contains the proper ultraviolet divergence which satisfies the unitary condition of Eq.(3), $\delta \bar{V}$ is the correct CKM counterterm at one-loop level. The explicit result is as follows

$$\delta \bar{V}_{ij} = \frac{1}{4} \sum_k (\delta \bar{Z}_{ki}^{uL*} - \delta \bar{Z}_{ik}^{uL}) V_{kj} + \frac{1}{4} \sum_k V_{ik} (\delta Z_{jk}^{dL*} - \delta Z_{kj}^{dL}) + \frac{1}{4} V_{ij} \delta \bar{Z}_{ii[l]m_{d,i} \rightarrow m_{d,j}}^{uL} + \frac{1}{4} V_{ij} \delta Z_{jj[l]m_{u,j} \rightarrow m_{u,i}}^{dL} - \frac{1}{4} \sum_{kl} V_{ik} V_{lk}^* V_{lj} (\delta \bar{Z}_{ll[l]m_{d,l} \rightarrow m_{d,k}}^{uL*} + \delta Z_{kk[l]m_{u,k} \rightarrow m_{u,l}}^{dL*}) \quad (14)$$

IV. RELATIONSHIP BETWEEN THE UNITARITY AND THE GAUGE INDEPENDENCE OF CKM MATRIX

It has been proven that using Nielsen identities [22] any physical parameter's counterterm must be gauge independent [23], So does the CKM counterterm [13]. It has been point out that in order to keep the gauge independence of the CKM matrix the CKM renormalization prescription must keep the unitarity of the CKM matrix [13,8]. But this conclusion is only a hypothesis. An explicit Proof has not been given. Here we will give an explicit and sufficient Proof by studying the relationship between the one-loop and two-loop CKM counterterms.

Here one will encounter one question: if the CKM counterterms in lower-loop levels which are gauge independent but don't satisfy the unitary condition have been obtained, will the CKM counterterms in higher-loop levels be gauge dependent or not? Our answer for this question is yes. This can be seen from the influence of the one-loop CKM counterterm to the two-loop CKM counterterm. In order to elaborate this problem clearly we express the amplitude of $W^+ \rightarrow u_i \bar{d}_j$ in the following form

$$T(V^0) = T(V + \delta V) = T(V) + T'(V) \delta V + \frac{1}{2} T''(V) (\delta V)^2 + \dots \quad (15)$$

where the superscript ' denotes the partial derivative with respect to the CKM matrix and the amplitude T doesn't contain the CKM counterterm. To two-loop level, this equation becomes

$$T_2(V^0) = T_2(V) + T_1'(V) \delta V_1 + \delta V_2 A_L \quad (16)$$

where the subscripts "1" and "2" denote the 1-loop and 2-loop results. Since $T_2(V^0)$ must be gauge independent, the gauge dependence of δV_2 is determined by the gauge dependence of $T_1'(V) \delta V_1$ and $T_2(V)$. Using the facts that F_R and $G_{L,R}$ are gauge independent and don't contain CKM matrix element and the terms in the first bracket of Eq.(4) are gauge independent [13], we get

$$T_1'(V) \delta V_1|_\xi = [-\frac{\delta V_{ij}}{2V_{ij}} (\sum_{k \neq i} \delta \bar{Z}_{ik}^{uL} V_{kj} + \sum_{k \neq j} V_{ik} \delta Z_{kj}^{dL}) + \frac{V_{ij}}{2} \sum_{k,l} (\frac{2}{g} \frac{d(\delta g)}{dV_{kl}} + \frac{d(\delta Z_W)}{dV_{kl}} + \frac{d(\delta \bar{Z}_{ii}^{uL})}{dV_{kl}} + \frac{d(\delta Z_{jj}^{dL})}{dV_{kl}}) \delta V_{kl} + \frac{1}{2} (\sum_{k \neq i} \delta \bar{Z}_{ik}^{uL} \delta V_{kj} + \sum_{l,m,k \neq i} \frac{d(\delta \bar{Z}_{ik}^{uL})}{dV_{lm}} \delta V_{lm} V_{kj} + \sum_{k \neq j} \delta V_{ik} \delta Z_{kj}^{dL} + \sum_{l,m,k \neq j} V_{ik} \frac{d(\delta Z_{kj}^{dL})}{dV_{lm}} \delta V_{lm})] A_L \quad (17)$$

where the subscript "1" of δV_1 has been omitted and the subscript " ξ " denotes the gauge dependent part of the quantity. Omitting the imaginary part of the quark's self energies, we obtain

$$\begin{aligned}
T'_1(V)\delta V_1|_\xi = & \frac{\alpha A_L}{32\pi M_W^2 s_W^2 m_{d,j}^2} \sum_{k,l} (\delta V_{il} V_{kl}^* + V_{il} \delta V_{kl}^*) V_{kj} [-\xi_W^2 M_W^4 \ln \frac{m_{u,k}^2}{M_W^2} + 2\xi_W m_{d,j}^2 M_W^2 \ln \frac{m_{u,k}^2}{M_W^2} - m_{u,k}^4 \ln \xi_W \\
& + 2m_{d,j}^2 m_{u,k}^2 \ln \xi_W - 2\sqrt{-\xi_W^2 M_W^4 + 2\xi_W m_{d,j}^2 M_W^2 + 2\xi_W m_{u,k}^2 M_W^2 - m_{d,j}^4 - m_{u,k}^4 + 2m_{d,j}^2 m_{u,k}^2} \\
& (\xi_W M_W^2 - m_{d,j}^2 + m_{u,k}^2) \arctan \frac{\sqrt{-\xi_W^2 M_W^4 + 2\xi_W m_{d,j}^2 M_W^2 + 2\xi_W m_{u,k}^2 M_W^2 - m_{d,j}^4 - m_{u,k}^4 + 2m_{d,j}^2 m_{u,k}^2}}{-\xi_W M_W^2 + m_{d,j}^2 - m_{u,k}^2}] + \\
& \frac{\alpha A_L}{32\pi M_W^2 s_W^2 m_{u,i}^2} \sum_{k,l} (V_{lk}^* \delta V_{lj} + \delta V_{lk}^* V_{lj}) V_{ik} [-\xi_W^2 M_W^4 \ln \frac{m_{d,k}^2}{M_W^2} + 2\xi_W m_{u,i}^2 M_W^2 \ln \frac{m_{d,k}^2}{M_W^2} - m_{d,k}^4 \ln \xi_W \\
& + 2m_{u,i}^2 m_{d,k}^2 \ln \xi_W - 2\sqrt{-\xi_W^2 M_W^4 + 2\xi_W m_{u,i}^2 M_W^2 + 2\xi_W m_{d,k}^2 M_W^2 - m_{u,i}^4 - m_{d,k}^4 + 2m_{u,i}^2 m_{d,k}^2} \\
& (\xi_W M_W^2 - m_{u,i}^2 + m_{d,k}^2) \arctan \frac{\sqrt{-\xi_W^2 M_W^4 + 2\xi_W m_{u,i}^2 M_W^2 + 2\xi_W m_{d,k}^2 M_W^2 - m_{u,i}^4 - m_{d,k}^4 + 2m_{u,i}^2 m_{d,k}^2}}{-\xi_W M_W^2 + m_{u,i}^2 - m_{d,k}^2}]
\end{aligned} \tag{18}$$

with ξ_W the W boson gauge parameter. It is easy to be seen that if δV_1 satisfies the unitary condition of Eq.(3), $T'_1(V)\delta V_1|_\xi$ will be zero. Thus $T_2(V)$ must be gauge independent because in this case δV_2 is gauge independent [23]. On the other hand, if δV_1 doesn't satisfy the unitary condition $T'_1(V)\delta V_1|_\xi \neq 0$, δV_2 must be gauge dependent in order to keep the amplitude $T_2(V^0)$ gauge independent.

Although this conclusion is only about the two-loop result, it is a sufficient condition to verify the proposition: only if the unitarity of the CKM matrix is kept in the CKM renormalization prescription, the renormalized CKM matrix and its counterterm are gauge independent.

V. CONCLUSION

In summary, We have investigated the present CKM matrix renormalization prescriptions and found Barroso's prescription [17] has a good idea. But we found it doesn't make the physical amplitude involving quark mixing ultraviolet finite. We then modified this prescription to solve this problem and have proven this revised prescription keep the unitarity of the CKM matrix at very high precision. So we can use this revised definition of the CKM counterterm in Eq.(9) in actual calculations. A more severe CKM renormalization prescription has been given in Ref. [8]. We also give an explicit Proof to prove that only if the CKM renormalization prescription keeps the unitarity of the CKM matrix the renormalized CKM matrix and its counterterm are gauge independent.

Acknowledgments

The author thanks professor Xiao-Yuan Li for his useful guidance and Dr. Hu qingyuan for his sincerely help (in my life). The author also thanks Prof. B. A. Kniehl very much for his kindly criticism to help him improving this work.

-
- [1] N.Cabibbo, Phys. Rev. Lett. **10**, 531 (1963);
M.Kobayashi and K.Maskawa, Prog. Theor. Phys. **49** (1973) 652.
 - [2] Particle Data Group, D.E. Groom et al., Eur. Phys. J. C**15** (2000) 1;
A. Höcker, H. Lacker, S. Laplace, F.Le Diberder, Eur. Phys. J. C**21** (2001) 225;
M. Beneke et al., hep-ph/0003033.
 - [3] L.-L. Chau and W.Y. Keung, Phys. Rev. Lett. **53** (1984) 1802;
J.D. Bjorken, Phys. Rev. D**39** (1989) 1396;
G.C. Branco and L. Lavoura, Phys. Lett B**208** (1988)123.
 - [4] A. Sirlin, Rev. Mod. Phys. **50** (1978) 573;
W.J. Marciano and A. Sirlin, Nucl. Phys B**93** (1975) 303.
 - [5] W.J. Marciano, A. Sirlin, Nucl. Phys. B**93** (1975) 303.
 - [6] A. Denner, T. Sack, Nucl. Phys. B**347** (1990) 203.
 - [7] J. Manzano, hep-ph/0208068;
D. Espriu, J. Manzano, P. Talavera, Phys. Rev. D**66** (2002) 076002;
A. Pilaftsis, Phys. Rev. D**65** (2002) 115013.

- [8] K.-P. O. Diener, B. A. Kniehl, Nucl. Phys. B**617** (2001) 291.
- [9] C. Becchi, A. Rouet, and R. Stora, Commun. Math. Phys. **42** (1975) 127; Ann. Phys. (N.Y.) **98** (1976) 287.
- [10] W.A. Bardeen, A.J. Buras, D.W. Duke, T. Muta, Phys. Rev. D**18** (1978) 3998.
- [11] D.J. Gross and F. Wilczek, Phys. Rev. D**8** (1973) 3633;
W.E. Caswell and F. Wilczek, Phys. Lett. B**49** (1974) 291;
H. Kluberg-Stern and J.B. Zuber, Phys. Rev. D**12** (1975) 467; Phys. Rev. D**12**, (1975) 482.
- [12] D. Espriu, J. Manzano, Phys. Rev D**63** 073008.
- [13] P. Gambino, P.A. Grassi, F. Madricardo, Phys. Lett. B 454 (1998) 98.
- [14] B.A. Kniehl, F. Madricardo, M. Steinhauser, Phys. Rev. D 62 (2000) 073010.
- [15] Y. Yamada, Phys. Rev. D**64** (2001) 036008.
- [16] J.M. Cornwall, Phys. Rev. D**26** (1982) 1453;
J.M. Cornwall and J. Papavassiliou, Phys. Rev. D**40** (1989) 3474;
J. Papavassiliou, Phys. Rev. D**41** (1990) 3179; Phys. Rev. D**51** (1995) 856;
G. Degrassi, B.A. Kniehl and A. Sirlin, Phys. Rev. D**48** (1993) 3963.
- [17] A. Barroso, L. Brucher, R. Santos, Phys. Rev. D 62 (2000) 096003.
- [18] Yong Zhou, J. Phys. G **29** (2003) 1.
- [19] T. Hahn, Comp. Phys. Comm. **140** (2001) 418.
- [20] R. Mertig, M. Bohm and A. Denner, Comp. Phys. Comm. **64** (1991) 345.
- [21] K. Fujikawa, B.W. Lee, A.I. Sanda, Phys. Rev. D**6** (1972) 2923.
- [22] N.K. Nielsen, Nucl. Phys. B**101** (1975) 173;
O. Piguet and K. Sibold, Nucl. Phys. B**253** (1985) 517;
P. Gambino and P.A. Grassi, Phys. Rev. D**62** (2000) 076002.
- [23] P. Gambino and P.A. Grassi, Phys. Rev. D**62** 076002.